


**BELEMNITE**


Pachyteuthis abbreviata
A squid relative from the Jurassic, *Pachyteuthis* had a calcite guard. It was about 10 cm (4 in) long.

CYSTOID

Pseudocrinites bifasciatus
Characterized by rhomboid-shaped respiratory structures, *Pseudocrinites* lived in Silurian and Devonian times. It was attached to the seabed by its stem.



respiratory structure

**DEVONIAN CRINOID**

Cupressocrinites crassus
Up to 3 cm (1¼ in) in diameter, this crinoid from Devonian strata had a tall, five-sided cup at the end of its stem.

flexible arm

stem



densely packed branches

JURASSIC CRINOID

Pentacrinites sp.
Named for its five-sided ossicles (stem segments), *Pentacrinites* grew to over 1 m (3¼ ft) in height. It is often found attached to fossil wood.

**BRITTLESTAR**

Lapworthura miltoni
An early fossil brittlestar, *Lapworthura* is from Ordovician and Silurian strata. It grew to 10 cm (4 in) in diameter. This species had five relatively short, thick arms.

**STARFISH**

Tropidaster pectinatus
This extinct starfish from the Early Jurassic was about 2.5 cm (1 in) wide. It had five thick arms.

BLASTOID

Pentremites pyriformis
This echinoderm from the blastoid group lived during Carboniferous times. It had long, arm-like structures that were used for feeding.

**SEA URCHIN**

Hemicidaris intermedia
This common sea urchin from Jurassic strata grew to around 4 cm (1½ in) in diameter. Its many bumps (tubercles) supported stout spines.

HEART URCHIN

Lovenia sp.
A heart-shaped, burrowing sea urchin, *Lovenia* is known from the Palaeocene and still exists today. It is up to 5 cm (2 in) in diameter.



tubercle where spine was attached