



SCOLIDOSAURUS FOOT
Scelidosaurus harrisonii

Found in Early Jurassic rocks, *Scelidosaurus* grew to 4 m (13 ft) in length and was covered with sharp, bony knobs. It had long toes and blunt claws.



STEGOSAURUS PLATE
Stegosaurus sp.

A Late Jurassic plant-eater, *Stegosaurus* grew up to 9 m (30 ft) in length. It had two rows of huge bony plates running along its back.




ANKYLOSAURUS SKULL
Ankylosaurus magniventris

Found in Cretaceous rocks, *Ankylosaurus* was a heavily armoured plant-eater. It grew to about 6 m (20 ft) in length.




EUOPLOCEPHALUS TAIL CLUB
Euoplocephalus tutus

Growing to 7 m (23 ft) in length, *Euoplocephalus* lived in the Late Cretaceous. A bony club at the tip of its tail was probably used for defence.



PARASAUROLOPHUS SKULL
Parasaurolophus walkeri

A Cretaceous herbivore, *Parasaurolophus* had a long, curved, hollow crest on its skull. This may have been used to make deep, resonating calls.



HYPsilOPHODON TOE
Hypsilophodon foxii

Dating back to the Cretaceous Period, *Hypsilophodon* was a swift-moving herbivore. It grew up to 2.3 m (7½ ft) in length.



PACHYCEPHALOSAURUS SKULL
Pachycephalosaurus wyomingensis

Living at the end of the Cretaceous Period, *Pachycephalosaurus* had a thick, domed skull and grew up to 5 m (16 ft) in length.



STEGOCERAS SKULL
Stegoceras validum

Found in Cretaceous strata, *Stegoceras* grew to 2 m (6½ ft) in length. Its small, serrated teeth indicate it was probably a herbivore.


TRICERATOPS SKULL
Triceratops prorsus

A massive horned and plated skull characterized *Triceratops*, a plant-eater from Late Cretaceous strata.



PSITTACOSAURUS SKELETON
Psittacosaurus sp.

One of the earliest horned dinosaurs, *Psittacosaurus* is found in Cretaceous strata. It was a plant-eater and grew to 2 m (6½ ft) in length.



toothless
beak



large nasal cavity

cheek horn

STYRACOSAURUS SKULL
Styracosaurus albertensis

Although similar to *Triceratops*, *Styracosaurus* had slender horns along the back of its skull. It is found in Late Cretaceous rocks.