

Unveiling the Deadly Nexus: Tobacco and Cancer

In the relentless battle against cancer, one adversary looms larger than most: tobacco. Decades of research have showed the intricate and devastating association between tobacco use and various forms of cancer. From lung cancer to pancreatic and beyond, the evidence linking tobacco consumption to cancer incidence and mortality is unequivocal.

The Lung Cancer Epidemic:

At the forefront of this deadly nexus is lung cancer, which is one of the most prevalent and lethal forms of cancer worldwide. Main causative agent is tobacco smoke which contains over 7,000 chemicals, including at least 70 known carcinogens. The inhalation of these toxic fumes wreaks havoc on the delicate lung tissue, leading to cellular mutations and the unchecked proliferation of cells.

According to the World Health Organization (WHO), tobacco use is responsible for approximately 90% of all lung cancer cases. Furthermore, research published in the Journal of the National Cancer Institute estimates that smokers are 15 to 30 times more likely to develop lung cancer compared to non-smokers.

Beyond the Lungs:

While lung cancer garners much of the attention in discussions about tobacco-related malignancies, the harmful effects of tobacco extend far beyond the respiratory system. Studies have implicated tobacco use in the development of cancers affecting the mouth, throat, esophagus, pancreas, bladder, kidney, cervix, and more.

For instance, a meta-analysis published in the Journal of the American Medical Association found that smoking is associated with a two-fold increase in the risk of pancreatic cancer. Similarly, the International Agency for Research on Cancer (IARC) has classified tobacco smoke as a known cause of bladder cancer.

The Role of Secondhand Smoke:

The peril of tobacco extends beyond the smoker themselves, enveloping innocent bystanders in its toxic embrace. Secondhand smoke, which contains many of the same carcinogens as direct smoke, poses a significant risk to non-smokers. The American Cancer Society reports that exposure to secondhand smoke



Dr. Itesh Khatwani
Consultant-Medical Oncologist
Ahmedabad

increases the risk of lung cancer by 20-30% in non-smoking adults and is a known cause of sudden infant death syndrome (SIDS) in infants.

Conclusion:

As we confront the scourge of cancer, it is imperative to acknowledge and address the deadly nexus between tobacco and malignancy. Public health initiatives aimed at tobacco control, including smoking cessation programs and tobacco taxation, are crucial steps in mitigating the devastating impact of tobacco-related cancers.

References:

1. World Health Organization (WHO). Tobacco and Cancer.
2. Journal of the National Cancer Institute. Tobacco Smoking and Cancer: A Meta-Analysis.
3. Journal of the American Medical Association. Cigarette Smoking and Pancreatic Cancer.
4. International Agency for Research on Cancer (IARC). IARC Monographs on the Evaluation of Carcinogenic Risks to Humans: Tobacco Smoke and Involuntary Smoking.
5. American Cancer Society. Secondhand Smoke.

Approval ID: IN-16533 Approval Date: 27/05/2025 Expiry Date: 27/05/2027

Disclaimer: This article is for general awareness of the people and should not be construed as treatment advice. Please consult your doctor in case you have any symptoms or require further information/clarification. **This public awareness initiative is supported by AstraZeneca Pharma India Limited.**