INDIAN GEOGRAPHY

27. Consider the map of India given below :



The map shows

(a) an isohyet
(b) allIndia water divide
(c) a combination of national highways
(d) a combination of major railway routes

Ans. (b) : The map shows all-India water divide. The major water divides of India are as follow:

Western Ghats:

The Western Ghats is a mountain range that runs parallel to the western coast of the Indian peninsula, located entirely in India. It forms a major water divide in the peninsula.

Aravali Range:

The Aravali Hills lie on the western and north-western margins of the peninsular plateau.

In this way, Aravali acts as water divide of India.

28. Which one of the following types of erosion is responsible for the formation of Chambal Ravines ?

(a) Splash (b) Sheet
(c) Rill (d) Gully

Ans. (d) : Badland topography is a major feature of the Chambal valley and it is characterized by an undulating flood plain, gullies and ravines. Ravines are a type of fluvial erosional feature and are formed as a result of constant vertical erosion by streams and rivers flowing over semiarid and arid regions.

The region through which the Chambal river flows does not receive enough rainfall to create ravines that are 60 – 80 m deep.

29. In ancient Indian historical geography the name 'Ratnakara' denoted

(a) the Arabian Sea
(b) the Bay of Bengal
(c) the Indian Ocean
(d) the confluence of the Ganga, the Yamuna and the mythical Saraswati at Prayaga

Ans. (c) : In ancient Indian historical geography the name 'Ratnakara' denoted the Indian ocean.

Ratnakara is the sea between India and Sri Lanka, which has been famous for beautiful gems especially pearls since ancient times. In 'Raghuvansh', the great poet Kalidas has used the word 'Ratnakara' for the sea situated between Indo-Srilanka.

30. The average elevation of Tibetan Plateau above sea level is

(a) 2 km (b) 3 km
(c) 4 km (d) 5 km

Ans. (d) : The Tibetan plateau is also called the plateau of Tibet or the inland plateau of Asia. It is the largest and highest plateau in the world with an average elevation of around 4,500 to 5,000 meters. The Tibetan plateau is known as "the roof of the world".

31. The only anthropoid ape to be found in India is the

(a) hanuman monkey
(b) liontailed macaque of Western Ghats
(c) slow loris of Assam
(d) whitebrowed gibbon of Assam

Ans. (d) : The only anthropoid ape to be found in India is the whitebrowed gibbon of Assam.

The western hoolock gibbon has a much wider range, as it found in all states of the north-east, restricted between the south of the Brahmaputra river and east of the Dibang river. The eastern hoolock gibbon inhabits specific pockets of Arunachal Pradesh and Assam in India.

32. The Himalayas are formed of parallel fold ranges, of which the oldest range is

(a) the Siwalik Range
(b) the Lesser Himalayas
(c) the Great Himalayan Range
(d) the Dhaula Dhar Range


Ans. (c) : The oldest range of Himalayas are the Great Himalayan Range. The Himalayan Ranges can be grouped into four parallel longitudinal Mountain belts of varying width, each having distinct physiographic features and its own geological history. They are designated, from south to north, as the outer, or sub-Himalayas (also called the Siwalik Range); the lesser, or lower, Himalayas; the Great Himalayan Range (Great Himalayas); and the Tethys, or Tibetan, Himalayas. Farther north lie the Trans-Himalayas in Tibet proper. From west to east the Himalayas are divided broadly into three mountainous regions: western, central, and eastern.

33. The hallmark of watershed development in the semiarid regions of India is the

(a) establishment of sheds on large scale, so that the rain water does not evaporate
(b) undertaking of earthworks, soil conservation measures and treeplantation, so as to conserve soil moisture and recharge underground water
(c) drilling deep tubewells so as to tap water at the rockstrata level
(d) setting up a system of tanks by banking water from seasonal rivers

Ans. (b) : The agriculture in low rainfall areas of eastern Rajasthan is characterized by high risks from drought degraded natural resources and pervasive poverty, food insecurity and mal-nutrition. In this region, water is the main limiting factor for upgrading rainfed agriculture. For such areas integrated watershed management is recognized as a potential approach for agriculture growth and rehabilitation of fragile and degraded lands.

34. Consider the map of India given below :



The map, read with the index provided, refers to the distribution of

(a) winter rains (b) pressure
(c) annual rainy days (d) mean temperature

Ans. (d) : The map, read with the index provided, refers to the distribution of mean temperature. India is a vast country and can be categorized into six main subcategories that have been described by the koppen climate classification. The annual temperature of the country can be classified according to the season that comprise of the summer season, winter season and the rainy season.

35. Assertion (A): The monsoonal rainfall decreases as one goes towards the west and northwest in the Ganga plain.

Reason (R): The moisture bearing monsoonal winds go higher up as one moves up in the Ganga plain.

In the context of the above two statements which one of the following is correct ?

(a) Both A and R are true, and R is the correct explanation of A
(b) Both A and R are true, but R is not a correct explanation of A
(c) A is true, but R is false
(d) A is false, but R is true

Ans. (a) : The monsoonal rainfall decreases as one goes towards the west and north-west in the Ganga plain due to the moisture bearing Monsoonal winds go higher up as one moves up in the Ganga plain.

36. If safe storage is to be ensured, the moisture content of foodgrains at the time of harvesting should not be higher than

(a) 14% (b) 16%
(c) 18% (d) 20%

Ans. (a) : If safe storage is to be ensured, the moisture content of food grains at the time of harvesting should not be higher than 14%. One of the most critical physiological factors in successful grains storage is the moisture content of the crop. High moisture content leads to storage problems because it encourages fungal and insect problems, respiration and germination. However, moisture content in the growing crop is naturally high and only starts to decrease as the crop reaches maturity and the grains are drying.

37. The highest milk producing breed of goat in India is

(a) Barbari (b) Jamunapari
(c) Black Bengal (d) Beetal

Ans. (b) : The highest milk producing breed of goat in Indian is Jamunapari. This goat breed is native to India subcontinent and the goat breed name is derived from the river "Jamuna Par" in Uttar Pradesh. Jamunapari goats are exported to Indonesia and they have great success over there. Basically this goat breed is reared for both milk and meat.

38. Which of the following fish are bred exclusively in the cold regions of India ?

I. Catla.
II. Tinca.
III. Bhekti.
IV. Trout.

Choose the correct answer from the codes given below.

Codes:

(a) I and lV (b) I and IV
(c) II and IV (d) III and IV