Note

The set of all points in the cartesian plane can be viewed as the set of all ordered pairs (x, y) where x, y are real numbers. In fact, $\mathbb{R}\times\mathbb{R}$ is the set of all points which we call as the cartesian plane.

Activity 1

Let $A = \{x \mid x \in \mathbb{N}, x \leq 4\}$, $B = \{y \mid y \in \mathbb{N}, y < 3\}$

Represent $A \times B$ and $B \times A$ in a graph sheet. Can you see the difference between $A \times B$ and $B \times A$?

Example 1.1 If $A = \{1,3,5\}$ and $B = \{2,3\}$ then (i) find $A \times B$ and $B \times A$. (ii) Is $A \times B = B \times A$? If not why? (iii) Show that $n(A\times B) = n(B\times A) = n(A)\times n(B)$

Solution Given that $A = \{1,3,5\}$ and $B = \{2,3\}$

(i) $A\times B = \{1,3,5\} \times \{2,3\} = \{(1,2), (1,3), (3,2), (3,3), (5,2), (5,3)\}$ ...(1)

$B \times A = \{2,3\} \times \{1,3,5\} = \{(2,1), (2,3), (2,5), (3,1), (3,3), (3,5)\}$ ...(2)

(ii) From (1) and (2) we conclude that $A \times B \neq B \times A$ as $(1,2) \neq (2,1)$ and $(1,3) \neq (3,1)$, etc.

(iii) $n(A)=3$; $n(B) = 2$.

From (1) and (2) we observe that, $n(A\times B) = n(B\times A) = 6$;

we see that, $n(A) \times n(B) = 3 \times 2 = 6$ and $n(B) \times n(A) = 2\times 3 = 6$

Hence, $n(A\times B) = n(B\times A) = n(A) \times n(B) = 6$.

Thus, $n(A\times B) = n(B\times A) = n(A) \times n(B)$.

Example 1.2 If $A\times B = \{(3,2), (3,4), (5,2), (5,4)\}$ then find A and B.

Solution $A\times B = \{(3,2), (3,4), (5,2), (5,4)\}$

We have A = {set of all first coordinates of elements of $A \times B$}. $\therefore$ $A = \{3,5\}$

B = {set of all second coordinates of elements of $A \times B$}. $\therefore$ $B = \{2,4\}$

Thus $A = \{3,5\}$ and $B = \{2,4\}$.

Example 1.3 Let $A = \{x \in \mathbb{N} \mid 1 < x < 4\}$, $B = \{x \in \mathbb{W} \mid 0 \leq x < 2\}$ and $C = \{x \in \mathbb{N} \mid x < 3\}$. Then verify that

(i) $A \times (B \cup C) = (A \times B) \cup (A \times C)$ (ii) $A \times (B \cap C) = (A \times B) \cap (A \times C)$

Solution $A = \{x \in \mathbb{N} \mid 1 < x < 4\} = \{2, 3\}$, $B = \{x \in \mathbb{W} \mid 0 \leq x < 2\} = \{0,1\}$,

$C = \{x \in \mathbb{N} \mid x < 3\} = \{1,2\}$

(i) $A \times (B \cup C) = (A \times B) \cup (A \times C)$

$B \cup C = \{0,1\} \cup \{1,2\} = \{0,1,2\}$

$A \times (B \cup C) = \{2,3\} \times \{0,1,2\} = \{(2,0),(2,1),(2,2),(3,0),(3,1),(3,2)\}$ ...(1)

$A \times B = \{2,3\} \times \{0,1\} = \{(2,0),(2,1),(3,0),(3,1)\}$

$A \times C = \{2,3\} \times \{1,2\} = \{(2,1),(2,2),(3,1),(3,2)\}$

1 Relation and Functions.indd 4 15-12-2020 02:50:02 PM