Progress Check

1. The sum of all the observations divided by number of observations is _____.
2. If the sum of 10 data values is 265 then their mean is _____.
3. If the sum and mean of a data are 407 and 11 respectively, then the number of observations in the data are _____.

8.2 Measures of Dispersion

The following data provide the runs scored by two batsmen in the last 10 matches.

Batsman A: 25, 20, 45, 93, 8, 14, 32, 87, 72, 4

Batsman B: 33, 50, 47, 38, 45, 40, 36, 48, 37, 26

$$\text{Mean of Batsman A} = \frac{25+20+45+93+8+14+32+87+72+4}{10} = 40$$

$$\text{Mean of Batsman B} = \frac{33+50+47+38+45+40+36+48+37+26}{10} = 40$$

The mean of both datas are same (40), but they differ significantly.




Fig. 8.1(a)




Fig. 8.1(b)

From the above diagrams, we see that runs of batsman B are grouped around the mean. But the runs of batsman A are scattered from 0 to 100, though they both have same mean.

Thus, some additional statistical information may be required to determine how the values are spread in data. For this, we shall discuss Measures of Dispersion.

Dispersion is a measure which gives an idea about the scatteredness of the values.

Measures of Variation (or) Dispersion of a data provide an idea of how observations spread out (or) scattered throughout the data.

Different Measures of Dispersion are

1. Range
2. Mean deviation
3. Quartile deviation
4. Standard deviation
5. Variance
6. Coefficient of Variation

8 Stats and Probability.indd 303 15-12-2020 03:24:05 PM