$$\frac{6}{6+3x} = 3 \times \frac{x}{6+3x}$$

$$3x = 6 \text{ gives, } x = 2$$

(i) Number of black balls $= 2 \times 2 = 4$

(ii) Total number of balls $= 6 + (3 \times 2) = 12$

Example 8.25 A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the numbers 1, 2, 3, ...12. What is the probability that it will point to (i) 7 (ii) a prime number (iii) a composite number?




Fig. 8.6

Solution Sample space $S = \{1,2,3,4,5,6,7,8,9,10,11,12\}$; $n(S) = 12$

(i) Let A be the event of resting in 7. $n(A)=1$

$$P(A) = \frac{n(A)}{n(S)} = \frac{1}{12}$$

(ii) Let B be the event that the arrow will come to rest in a prime number.

$$B = \{2,3,5,7,11\};\ n(B) = 5$$

$$P(B) = \frac{n(B)}{n(S)} = \frac{5}{12}$$

(iii) Let C be the event that arrow will come to rest in a composite number.

$$C = \{4,6,8,9,10,12\};\ n(C) = 6$$

$$P(C) = \frac{n(C)}{n(S)} = \frac{6}{12} = \frac{1}{2}$$



Thinking Corner

What is the complement event of an impossible event?



Exercise 8.3

1. Write the sample space for tossing three coins using tree diagram.
2. Write the sample space for selecting two balls at a time from a bag containing 6 balls numbered 1 to 6 (using tree diagram).
3. If A is an event of a random experiment such that $P(A) : P(\bar{A}) = 17:15$ and $n(S)=640$ then find (i) $P(\bar{A})$ (ii) $n(A)$.
4. A coin is tossed thrice. What is the probability of getting two consecutive tails?
5. At a fete, cards bearing numbers 1 to 1000, one number on one card are put in a box. Each player selects one card at random and that card is not replaced. If the selected card has a perfect square number greater than 500, the player wins a prize. What is the probability that (i) the first player wins a prize (ii) the second player wins a prize, if the first has won?
6. A bag contains 12 blue balls and x red balls. If one ball is drawn at random (i) what is the probability that it will be a red ball? (ii) If 8 more red balls are put in the bag, and if the probability of drawing a red ball will be twice that of the probability in (i), then find x.
7. Two unbiased dice are rolled once. Find the probability of getting

   (i) a doublet (equal numbers on both dice) (ii) the product as a prime number
   (iii) the sum as a prime number (iv) the sum as 1

8 Stats and Probability.indd 323 15-12-2020 03:24:33 PM