

Fig 10.8: Kwashiorkor



Fig 10.9: Marasmus

8) Dehydration
9) Body temperature is sub-normal
10) Failure to thrive
11) Wrinkled skin - Old man's face
12) Grossly underweight

III. Marasmic Kwashiorkor

Children suffering from this disease show signs of both kwashiorkor and marasmus.

10.7.5 Differences between Kwashiorkor and Marasmus

Table 10.5: Differences between Kwashiorkor and Marasmus

Kwashiorkor	Marasmus
It develops in children whose diets are deficient of protein.	It is due to deficiency of proteins and calories.
It occurs in children between 6 months and 3 years of age.	It is common in infants under 1 year of age.
Subcutaneous fat is preserved.	Subcutaneous fat is not preserved
Oedema is present.	Oedema is absent.
Enlarged fatty liver.	No fatty liver.
Ribs are not very prominent.	Ribs become very prominent.
Lethargic	Alert and irritable.
Mild or no Muscle wasting.	Severe muscle wasting
Poor appetite.	Voracious feeder.
The person suffering from kwashiorkor needs adequate amounts of proteins.	The person suffering from marasmus needs adequate amount of proteins, fats and carbohydrates.

UNIT_10_149-173.indd 158 13-02-2020 19:27:45