ACTIVITY - 3

To compare quality characteristics of fresh and deteriorated raw eggs

Materials Needed:

1 fresh egg

2 deteriorated egg (held at room temperature for at least two weeks)

Procedure

1. Break one fresh egg out of the shell onto a flat plate. Be careful not to damage the egg.
2. Keep the shell. Inspect the air cell in the large blunt end of the egg shell.
3. Observe the thick and thin egg white: the height, diameter, and the color of the yolk; the position of the yolk in the white; the chalazae and record the observations.
4. Repeat steps 1-3 using a deteriorated egg.


4.4.4 The properties of egg

There are three main properties of proteins in eggs which enable them to be used in so many different ways in cookery.


1. Egg proteins coagulate on heating.
2. Egg proteins stretch when beaten and hold air in the structure.
3. Egg yolk proteins are good emulsifying agents.

4.4.5 Uses of egg in cookery

Eggs can be used in many ways in cookery. Eggs when used alone or in combination with other foods they become the major protein source of a meal. Eggs can be used as boiled, scrambled, fried (omelettes) or poached for table use. Eggs are used as :



In a good quality egg the white is thick and stands high. Yolk is firm, round and high.
In a poor quality egg the yolk is somewhat flattened and enlarged. The white appears watery.



When the egg sinks in water it is considered as fresh.

Poor quality eggs float due to increase in size of the air cell and due to loss of moisture.

- **Thickening Agent:** Egg proteins coagulate on heating. Therefore, eggs can be used as thickening agents for making stirred and baked custards, soups and puddings.
- **Binding Agent:** Egg proteins coagulate between 65 and 70°C and

UNIT-04_055-073.indd 65 22-01-2020 12:47:00