

above the plane of the paper and the dashed wedge (·····) (or a dashed line) is used to depict the bond below the plane. The bonds lying in the plane of the paper are shown by normal lines.




Fisher projection formula:

This is a method of representing three dimensional structures in two dimension. In this method, the chiral atom(s) lies in the plane of paper. The horizontal substituents are pointing towards the observer and the vertical substituents are away from the observer. Fisher projection formula for tartaric acid is given below.




Sawhorse projection formula:

Here the bond between two carbon atoms is drawn diagonally and slightly elongated. The lower left hand carbon is considered lying towards the front and the upper right hand carbon towards the back. The Fischer projection inadequately portrays the spatial relationship between ligands attached to adjacent atoms. The sawhorse projection attempts to clarify the relative location of the groups.



Newman projection formula:

In this method the molecules are viewed from the front along the carbon-carbon bond axis. The two carbon atom forming the σ bond is represented by two circles. One behind the other so that only the front carbon is seen. The front carbon atom is shown by a point where as the carbon lying further from the eye is represented by the origin of the circle. Therefore, the C-H bonds of the front carbon are depicted from the circle while C-H bonds of the back carbon are drawn from the circumference of the circle with an angle of 120^0 to each other.



11.5. Isomerism in organic compounds:

The term 'isomerism' was given by Berzelius, and its represents of existence

Unit 11 revised - jagan.indd 131 22-01-2021 12:42:25