Steroisomerism:

11.5.3 Geometrical isomerism:

Geometrical isomers are the stereoisomers which have different arrangement of groups or atoms around a rigid frame work of double bonds. This type of isomerism occurs due to restricted rotation of double bonds, or about single bonds in cyclic compounds.

In alkenes, the carbon-carbon double bond is sp^2 hybridized. The carbon-carbon double bond consists of a σ bond and a π bond. The σ bond is formed by the head on overlap of sp^2 hybrid orbitals. The π bond is formed by the side wise overlap of 'p' orbitals. The presence of the π bond lock the molecule in one position. Hence, rotation around C=C bond is not possible. This restriction of rotation about C-C double bond is responsible for geometrical isomerism in alkenes.



cis - 2-butene *Trans* 2-butene

These two compounds are termed as geometrical isomers and are distinguished from each other by the terms cis and trans. The cis isomer is one in which two similar groups are on the same side of the double bond. The trans isomers is that in which the two similar groups are on the opposite side of the double bond, hence this type of isomerism is often called cis-trans isomerism.

The cis-isomer can be converted to trans isomer or vice versa is only if either isomer is heated to a high temperature or absorbs light. The heat supplies the energy (about 62kcal/mole) to break the π bond so that rotation about σ bond becomes possible. Upon cooling, the reformation of the π bond can take place in two ways giving a mixture both cis and trans forms of trans-2-butene and cis-2-butene.

Unit 11 revised - jagan.indd 135 22-01-2021 12:42:26