Carbon combines with chlorine to form carbon tetrachloride, which is a liquid and insoluble (immiscible) in water. Sodium combines with chlorine atom to form sodium chloride, a hard and brittle compound that readily dissolves in water. The possible reason for these observations lies in the type of interaction that exists between the atoms of these molecules and these interactions are responsible for holding the atoms/ions together. The interatomic attractive forces which hold the constituent atoms/ions together in a molecule are called chemical bonds.

Why do atoms combine only in certain combinations to form molecules? For example oxygen combines with hydrogen to give water (H_2O) and with carbon it gives carbon dioxide (CO_2). The structure of water is 'V' shaped while that of the carbon dioxide is linear. Such questions can be answered using the principles of chemical bonding. In this unit we will analyse the various theories and their principles, which were developed over the years to explain the nature of chemical bonding.

10.1.1 Kossel – Lewis approach to chemical bonding

A logical explanation for chemical bonding was provided by Kossel and Lewis in 1916. Their approach to chemical bonding is based on the inertness of the noble gases which have little or no tendency to combine with other atoms. They proposed that the noble gases are stable due to their completely filled outer shell electronic configuration. Elements other than noble gases, try to attain the completely filled electronic configurations by losing, gaining or sharing one or more electrons from their outer shell. For example, sodium loses one electron to form Na^+ ion and chlorine accepts that electron to give chloride ion (Cl^-), enabling both atoms to attain the nearest noble gas configuration. The resultant ions, Na^+ and Cl^- are held together by electrostatic attractive forces and the attractive force is called a chemical bond, more specifically an electrovalent bond.

$$Na \longrightarrow Na^+ + e^-$$
$$\mathbf{[Ne]\ 3s^1} \qquad\qquad \mathbf{[Ne]}$$

$$Cl + e^- \longrightarrow Cl^-$$
$$\mathbf{[Ne]\ 3s^2\ 3p^5} \qquad\qquad \mathbf{[Ar]}$$

$$Na^+ + Cl^- \longrightarrow NaCl$$

G. N. Lewis proposed that the attainment of stable electronic configuration in molecules such as diatomic nitrogen, oxygen etc… is achieved by mutual sharing of the electrons. He introduced a simple scheme to represent the chemical bond and the electrons present in the outer shell of an atom, called Lewis dot structure. In this scheme, the valence electrons (outer shell electrons) of an element are represented as small dots around the symbol of the element. The first four valence electrons are denoted as single dots around the four sides of the atomic symbol and then the fifth onwards, the electrons are denoted as pairs. For example, the electronic configuration of nitrogen is $1s^2$, $\mathbf{2s^2, 2p^3}$. It has 5 electrons in its outer shell (valence shell). The Lewis structure of nitrogen is as follows.



Fig 10.1 Lewis Structure of Nitrogen atom

Similarly, Lewis dot structure of carbon, oxygen can be drawn as shown below.

Unit 10 layout - jagan.indd 68 22-01-2021 12:33:01