10.5.5 Resonance

When we write Lewis structures for a molecule, more than one valid Lewis structures are possible in certain cases. For example let us consider the Lewis structure of carbonate ion $[CO_3]^{2-}$.

The skeletal structure of carbonate ion (The oxygen atoms are denoted as O_A, O_B & O_C



Total number of valence electrons = [1 x 4(carbon)] + [3 x 6 (oxygen)] + [2 (charge)] = 24 electrons.

Distribution of these valence electrons gives us the following structure.



Complete the octet for carbon by moving a lone pair from one of the oxygens (O_A) and write the charge of the ion (2-) on the upper right side as shown in the figure.



Fig 10. 14 (a) Lewis Structure of CO_3^{2-}

In this case, we can draw two additional Lewis structures by moving the lone pairs from the other two oxygens (O_B and O_C) thus creating three similar structures as shown below in which the relative position of the atoms are same. They only differ in the position of bonding and lone pair of electrons. Such structures are called resonance structures (canonical structures) and this phenomenon is called resonance.



Fig 10. 14 (b) Resonance structures of CO_3^{2-}

It is evident from the experimental results that all carbon-oxygen bonds in carbonate ion are equivalent. The actual structure of the molecules is said to be the resonance hybrid, an average of these three resonance forms. It is important to note that carbonate ion does not change from one structure to another and vice versa. It is not possible to picturise the resonance hybrid by drawing a single Lewis structure. However, the following structure gives a qualitative idea about the correct structure.

Unit 10 layout - jagan.indd 78 22-01-2021 12:33:09