
Bonding in Ethylene:

The bonding in ethylene can be explained using hybridisation concept. The molecular formula of ethylene is C_2H_4. The valency of carbon is 4. The electronic configuration of valence shell of carbon in ground state is $[He]2s^2\ 2p_x^1\ 2p_y^1\ 2p_z^0$. To satisfy the valency of carbon promote an electron from 2s orbital to $2p_z$ orbital in the excited state.



In ethylene both the carbon atoms undergoes sp^2 hybridisation involving 2s, $2p_x$ and $2p_y$ orbitals, resulting in three equivalent sp^2 hybridised orbitals lying in the xy plane at an angle of 120^0 to each other. The unhybridised $2p_z$ orbital lies perpendicular to the xy plane.

Formation of sigma bond:

One of the sp^2 hybridised orbitals of each carbon lying on the molecular axis (x-axis) linearly overlaps with each other resulting in the formation a C-C sigma bond. Other two sp^2 hybridised orbitals of both carbons linearly overlap with the four 1s orbitals of four hydrogen atoms leading to the formation of two C-H sigma bonds on each carbon.

Formation of Pi (π) bond:

The unhybridised $2p_z$ orbital of both carbon atoms can overlap only sideways as they are not in the molecular axis. This lateral overlap results in the formation a pi(π) bond between the two carbon atoms as shown in the figure.




Fig 10.27 sp^2 Hybridisation : C_2H_4

Unit 10 layout - jagan.indd 95 22-01-2021 12:33:29