Objective Type Questions

7. The ratio of ₹1 to 20 paise is _____.

   (a) 1 : 5 (b) 1 : 2 (c) 2 : 1 (d) 5 : 1

8. The ratio of 1 l to 50 ml is _____.

   (a) 1 : 5 (b) 1 : 20 (c) 20 : 1 (d) 5 : 1

9. The length and breadth of a window are in 1m and 70 cm respectively. The ratio of the length to the breadth is _____.

   (a) 1 : 7 (b) 7 : 1 (c) 7 : 10 (d) 10 : 7

10. The ratio of the number of sides of a triangle to the number of sides of a rectangle is


    (a) 4 : 3 (b) 3 : 4 (c) 3 : 5 (d) 3 : 2

11. If Azhagan is 50 years old and his son is 10 years old then the simplest ratio between the age of Azhagan to his son is

    (a) 10 : 50 (b) 50 : 10 (c) 5 : 1 (d) 1 : 5

3.2.5 Equivalent Ratios

We can get equivalent ratios by multiplying or dividing the numerator and denominator by a common number. This is clear from the following example. Let us find the ratio between breadth and length of the following rectangles given below.



- Ratio of breadth to length of **rectangle A** is **1 : 2** (already in simplest form)
- Ratio of breadth to length of **rectangle B** is **2 : 4** (simplest form is 1 : 2)
- Ratio of breadth to length of **rectangle C** is **4 : 8** (simplest form is 1 : 2)
- Thus, the ratios of breadth and length of rectangles **A**, **B** and **C** are said to be *equivalent ratios*.
- That is, the ratios **1 : 2** = **2 : 4** = **4 : 8** are *equivalent*.

6th_Chapter 3_055-075.indd 52 1/27/2020 1:45:27 PM