
atmosphere and they cause chemical reactions that break down **ozone** molecules and reduce the concentration of them. Nitrogen oxide released by emitted by supersonic aircrafts can also destroy the ozone molecules to break down. Ozone-depleting substances are present throughout the stratospheric ozone layer because they are transported great distances by atmospheric air motions. The severe depletion of the Antarctic ozone layer known as the "ozone hole" occurs because of the special atmospheric and chemical conditions that exist there and nowhere else on the globe. The very low winter temperatures in the Antarctic stratosphere cause polar stratospheric clouds (PSCs) to form. Special reactions that occur on PSCs, combined with the relative isolation of polar stratospheric air, allow chlorine and bromine reactions to produce the ozone hole in Antarctic springtime.

Satellite images of the earth over last decades observed that the atmospheric ozone layer is getting thinner. On October 2, 2015, the ozone hole was recorded to its maximum size of **28.2 million sq.km** over Antarctica (Figure 6.5). The size of the ozone hole is larger than the size of continent of North America. The ozone holes over Antarctica allow the ultraviolet radiation to enter and cause of global warming, skin cancer, eye cataract and even blindness.

Depletion of the **ozone layer** has consequences on human, animal, plants and micro organisms. This typically results from higher UV levels reaching us on earth. Research confirms that high levels of UV rays cause non-melanoma skin cancer.



TN_Govt_Geography_XI_Std Unit_06.indd 142 2/10/2020 6:00:15 PM