Fixing:

After smear preparation the glass slide should show for 2 to 3 times in a flame. Due to flaming the specimen get fixed perfectly on a slide and also some chemicals like formalin, Methyl alcohol, Mercuric chloride is used for fixing the specimen.

Stains: - Stains is used to make cellular shapes and arrangements visible. For e.g. The stains used in gram staining – **Crystal violet, saffranin**)

(decolorizer -Ethyl alcohol, Mordant agent – grams Iodine).



Ear wax having anti-microbial properties that reduce the feasibility of bacteria and fungus in the human ear.


Hanging Drop Method: - (Glass slit method).


In hanging drop method, a drop of culture is made to hang between glass slide and slit and viewed under microscope.

The advantage of hanging drop method is we can identify mobile bacteria. Some bacteria have flagella for motility. e.g. Monotrichous, Peritrichous flagella (e.g. Proteus)

By cultural characteristics: -

Here bacteria are identified as group or culture as a whole and note individual bacteria some most bacteria grow in colonies and also divide fast. They can be easily grown into a culture in suitable nutrition media. Based on the characteristics of culture they can be identified as the size of colonies, type of elevation, margins, surface of colony, colour of culture.






Based on Antibiotic Resistance

Antibiotics (Ex-Penicillin) is added to the culture and measuring the resistance of microbe. The zone of inhibition surrounding the antiobiotic disc indicating sensitivity

No zone of inhibition surrounding the antibiotic disc indicating resistance.



By Biochemical test: -

Sugar fermentation test

- Litmus milk test
- Indole production test

Nursing-Gen_Chapter 06.indd 94 07/05/18 12:51 PM