
- Methyl Red test
- Citrate utilization test

By differential staining: -

The identification depends on staining of bacteria. And most bacteria can be stained by specific stain like crystal violet. Gram positive bacteria are stained by gram stain while Gram negative bacteria don't take up gram stain.

Mycobacterium tuberculosis bacteria can be stained by acid fast staining method.

Serological methods: - Here identification of bacteria is done by use of antibodies and antigens which are specific against the suspected bacteria. Antigens and antibodies are very specific and bind to single type of bacteria.



TEST

- VDRL, ELISA- Screening Test
- Western Blot - Confirmatory test

} Identification for HIV

Identification of bacteria is necessary to

I. Identify the disease
II. Select suitable drug
III. Evaluation of treatment progress
IV. For industrial purpose.
V. Storage

6.3.5 Types of specimen collection

Swabs: - It is usually collected in a sterile cotton swab, care should be taken so as to prevent contaminations of specimen. (e.g. Throat swab, Eye, Ear, Nose, Mouth, Vaginal, Abscesses swab). Materials should be taken only from the infected area.

Sputum: - It should be collected in a sterile container having wide mouth. Sputum should be collected directly after a cough and sent immediately to the laboratory.

Urine: - Urine specimen remains an important tool for clinical diagnosis. A correct urine result is influenced by the collection method, timing and handling (first morning sample, random sample). It should be collected in a sterile container.

Faeces: - Fresh stool should be collected for bacteriological examination. Specimen should be well covered and labelled.

For culture and parasite examination the specimen must be returned to the laboratory within one hour of collection.

Blood: - It is important that specimens are properly collected, prepared and preserved. When assisting the physician should adopt aseptic precautions so as to avoid contamination of specimen.

Always collect the blood specimen in hygienic area. Blood is carefully transfer from the syringe to the tube

Nursing-Gen_Chapter 06.indd 95 07/05/18 12:51 PM