iii) **Portal of exit:** -

- Pathogens have preferred portals of entry, they also have definite portal of exit.
- 3 common portal of exit.

  Respiratory tract – coughing, sneezing

  Gastro intestinal tract – Saliva, feaces

  Genital tract – Vagina and penis
- Arthropods and syringes provide a portal of exit for microbes in blood.

iv) **Modes of transmission**: -

Mode of transmission is through direct or indirect contact, droplet infection, vehicle transmission, air borne, arthropod vector.

v) **Portal of entry**: -

Organism can enter the body through skin, mucous membrane, gastro intestinal tract, intestinal tract, blood, genital tract.

vi. **Compromised host:** -

When a person acquires an infection depends on the susceptibility of an infectious agent.

- Individual degree of resistance to a pathogen.
- Patients with burn, surgical wounds, and suppressed immune system are the most susceptible.
- According to the virulence of the microbes.

ROUTES AND MEANS	TRANSMISSION	EXAMPLE (ORGANISM)
DIRECT CONTACT	PHYSICAL CONTACT	CHICKEN POX, MEASLES, RUBELLA
INDIRECT CONTACT	FOMITES (IN ANIMATE OBJECTS)	INFLUEZA (COMMON COLD)
DROPLET INFECTION	SALIVA, MUCUS	MYCOBACTERIUM TUBERCULOSIS
VEHICLE BORNE INFECTION	WATER, FOOD, AIR, BLOOD	V.CHOLERAE, SALMONELLA SHIGELLA
AIR BORNE INFECTION	WATER DROPLETS	INFLUENZA
VECTOR :-		
MECHANICAL & BIOLOGICAL	FLIES MOSQUITO LOUSE	SALMONELLA ADES EGYPTIA RICKETTSIA TYPI

Nursing-Gen_Chapter 06.indd 99 07/05/18 12:51 PM