Tears contain Bactericides such as lysozyme. So it kills bacteria (sometimes crying is good)

6.5.2 Course of Infection

Once a microorganism does overcome the defenses of the host, development of the disease follows a certain sequence of steps that tends to be similar whether the disease be acute or chronic.

- Incubation period
- Proximal stage
- Period of illness stage
- Period of decline
- Period of convalescence

Incubation Period: - The period of incubation is the time interval between the actual infection and the first appearance of signs and symptoms interval between entrance of pathogen into body and appearance of first symptoms.

e.g. Common cold - 1 to 2 days

Mumps - 18 days

Prodromal Stage: -The prodromal period is characterized by the appearance of the first mild signs and symptoms (low-grade, fever, fatigue). During this time microorganism grow and multiply and patient is more capable of spreading disease to others.

Illness Stage: - During this period of illness, the disease is at its height and all disease signs and symptoms are apparent.

e.g. Common cold - Sore throat, sinus

Mumps - Earache, high fever, parotid, salivary gland swelling


The severity of the patients' illness depends on the extent of infection, the pathogenicity of the microorganism, susceptibility of individuals.


Decline Stage: -

During the period of decline, the signs and symptoms subside.

Convalescence Period: -

During the period of convalescence, the body returns to its predeceased state, the health is restored. Length of recovery depends on severity of infection and patients' general health status. Recovery may take several days to month.





Nursing-Gen_Chapter 06.indd 100 07/05/18 12:51 PM