6. Hydrogen peroxide: 1-5% of solution is used in cleaning wounds and to remove pus from infected ears. Hydrogen peroxide is also used to clean the mouth. It is an oxidizing agent.
7. Potassium Permanganate: - It is an oxidizing agent used for cleaning the mouth with 1:1000 strength. It is also used for irrigation of wounds.
8. Carbolic acid (Phenol): - It is a good designating for feces, pus, blood and sputum. It is a skin irritant and a poison. Dissolves early in hot water. For thermometer 1:20 solution for a duration of 10 minutes.
9. Lysol: This is a phenol preparation mixed with soap. It is less poisonous than a carbolic acid but has a greater bactericidal action. 2% of solution for 6-8 hours is wed for disengaging livens.
10. Ethyl Alcohol: 70% is effective for skin disinfection. Certain gases like formaldehyde and glutaral-dehyde are used in disinfection of rooms.

6.6.1.4 Radiation*:*

The effects of radiation depend on its wave length, intensity, duration.

Ionizing radiation –

- Gamma rays and x-rays are both types of high energy electron (high frequency) electromagnetic radiation
- These rays can cause destruction of the DNA in microbes.
- The principal effect of ionizing radiation is the ionization of water, which forms highly reactive hydroxyl radicals.

Non-Ionizing radiation: UV rays

UV light damages the DNA of the exposed cells. It causes bonds to form between adjacent thymine dimers in DNA chain and inhibit replication.

USES*: -*

The radiations are used for sterilizing pharmaceutical and medical dental carries. (cold sterilization)

- Practical application is the UV lamp (germicidal property) in the microbiological laboratories.

6.6.1.5 Fumigation (0r) Gas Sterilization

Fumigation is a process of gaseous sterilization which is used for killing of microorganisms and prevention of microbial growth in air, surface of wall or floor.

- It is generally used in the pharmaceutical, operation theatres. Hospitals, and offices.
- For effective fumigation process is done according to the density. Humidity 60% and temperature never below 18° C in opened area around at a time of 1 hour to o16 hours it may be differ the gas kill all the spores, vegetative cells etc.,

Nursing-Gen_Chapter 06.indd 107 07/05/18 12:51 PM