4. If there is any discharge from the ear canal, inform it to the doctor.
5. Instruct the patient not to put any sharp objects like hair pin inside the ear.
6. In case of using hearing aid routine cleaning and proper insertion techniques should be followed.

7.11 CARE OF PATIENT WITH RETENTION OF URINE

Acute urinary retention is a medical emergency that requires prompt recognition and bladder drainage.



Shy bladder syndrome is a type of phobia were people are unable to pass urine in the public place

Nursing Management

1. Instruct the patient to warm up before attempting to urinate
2. Avoid alcohol intake
3. Advise the patient to drink a cup of coffee or tea to create urinary urgency.
4. Sitting in a tub of warm water or warm shower may help him to urinate.
5. Make the patient to scheduled toileting position to pass urine.
6. Running water sound may increase the urgency. Take him to the bathroom and hear the sound of running water.
7. Use a catheter with a retention balloon to empty the bladder.

7.12 Offering and removing the Bed Pan

Bed patients usually need to use a bedpan about once or three times a day. The ward may be closed with a screen for this purpose as a routine. However if a patient make request for a bedpan at another time, you should meet the request with understanding and without delay.

Method of giving a bed pan

1. Screen the bed.
2. Bring a covered bedpan to the beside and place it on the stool. The bed pan should be clear and dry. In cold weather, warm it first with hot water. Pad the seat of the bedpan, if the patient is very weak and emaciated.
3. Protect the bed with a rubber sheet if necessary.
4. Place the bedpan on the bedside for the patient.
5. Place your left hand beneath the lower back to aid the patient in raising the buttocks and place the bedpan in position without force. Adjust the bedpan comfortably for the patient. Cover and leave him alone unless he is too ill or weak
6. Get a toilet tray ready and bring it to the bedside contents of the tray are.

Sl. No.	Items
1	Jug with warm water
2	Soap
3	Wash cloth and towel
4	Bowl with rag pieces or cotton
5	Long artery forceps
6	Kidney tray and paper bag

Nursing-Gen_Chapter 07.indd 133 07/05/18 12:54 PM