To remove the empty bedpan

1. Let the patent wash himself if he is able to do so. Help him pouring water over the genitals, Remove the bedpan and give the patent water and soap for hand washing.
2. If the patient is helpless, use moistened rag pieces or cotton and the artery forceps and clean from front to back to prevent infection.

   Turn him on his side while removing the bedpan (a second person may be needed to help)
3. Make the patient comfortable.
4. Cover the bedpan, take it to the toilet room.
5. Observe the contents.
   - **A.** Proper position reduces patient's back strain.
   - **B.** Improper position of patient.





6. Empty the content and since the bed pan with cold water. Then clean it with the brush kept in soap solution the bed pan may be soaked in disinfectants. For one hour or sterilized if there are facilities for doing so.
7. Remove other articles from the bedside. Clean them and put them back in their proper place.
8. Wash your hands well.

Remove the screens, and leave the unit tidy. Record the time and observations.

7.13 Intake and Output Chart



Excessive sodium in the body retains water in the body and make your brain swell.

Intake and output records provide valuable information about fluid and electrolyte problems. As accurately recorded intake and output will identify sources of excessive intake of fluid losses. Intake should include Oral, IV, and tube Feedings and retained irrigations. Output includes urine, excess perspiration wound or tube drainage vomits and diarrhea. Estimate fluid loss from wounds and perspiration. Note the amount and colour of the urine.

Nursing-Gen_Chapter 07.indd 134 07/05/18 12:54 PM