3. Maintain an un -obstructed flow of urine. Empty the collecting bag regularly, and keep it below the level of the bladder.
4. Provide perineal care once or twice a day. Do not use lotion or powder near the catheter.
5. With long term use of a catheter, a long bag may be used.
6. Another catheter using some type of securement device to prevent catheter moving and urethral tension.
7. Use sterile technique whenever the system is open. e.g. sample collection for culture.

SUMMARY

- Hygiene is the practice of keeping yourself and your surroundings clean especially in order to prevent illness or the spread of disease.
- Factors affecting personal hygiene are physical, psychological and socio economic factors.
- Bed making contributes materially to the patient's comfort.
- There are too types. They are ordinary and special beds.
- Personal hygiene is one of the most effective ways to prevent the development and spread of infection.
- Bed bath is bathing a patient while he is bed.
- Routine care of shoulder and back is necessary to prevent bed sore.
- Foot and nail problems are due to improper trimming of nails, exposure to chemical and poorly fitted shoes.
- Good oral hygiene involves cleanliness, comfort and the moisturing the mouth structures and also prevent oral diseases.
- Hair wash is necessary for a patient to give comfort and sense of well being.
- Primary eye and ear care are the vital component of promotion of health.
- Acute urinary retension requires prompt recognition and bladder drainage.
- Catheterization is a routine medical procedure to drain out urine from the bladder.
- Nurse should meet the patient request of a bed pan with understanding and without delay.

Nursing-Gen_Chapter 07.indd 136 07/05/18 12:54 PM