

By volume, dry **air** contains 78.09% nitrogen, 20.95% oxygen, 0.93% argon, 0.04% carbon dioxide, and small amounts of other gases. **Air** also contains a variable amount of water vapor, on average around 1% at sea level, and 0.4% over the entire **atmosphere**.

Articles

1. Oxygen source
2. Mask (simple / or with venture adaptor high flow device of appropriate size)
3. Humidifier with distilled water
4. Flow meter
5. Gauze pieces
6. "No Smoking" sign.

Procedure

Sl.No	NURSING ACTION	RATIONALE
1	Determine need for oxygen therapy, Check physician's order for rate, device to be used and the concentration.	Reduces risk of error in administration.
2	Perform an assessment of vital signs, level of consciousness, lab values, etc. and record.	Provides a baseline for future assessment.
3	Assess risk factors of oxygen administration in patient and environment-like hypoxia drive in patients and faulty electrical connection.	Reduces risk of danger caused to patient. Oxygen is a combustible gas. Hypoxia drive in patients is essential to maintain respiration.
4	Explain procedure to patient and relatives and emphasize how he has to cooperate.	Reduces anxiety and enhances cooperation
5	Post "No Smoking" signals on the patient's door in view of patient and visitors and explain to them the dangers of smoking when oxygen is on flow.	Oxygen supports combustion; smoking in oxygen area can lead to fire hazards.
6	Wash hands	Reduces risk of transmission of microorganisms.
7	Set up oxygen equipments and humidifiers. a. Fill humidifier up to the level mark on it. b. Attach flow meter to source, set flow meter in 'off' position. c. Attach humidifier to base of flow meter d. Attach tubing and face mask to humidifier (if venture device is used attach the colour coded venture adapter to mask as appropriate) e. Regulate flow meter to prescribed level	Filling humidifier above this level will cause water to enter into tubing. Flow meter helps in monitoring and regulating oxygen flow to patient. Humidification helps to prevent drying of mucous membranes and promotes comfort of patient. Oxygen is a drug and is dangerous to administer at flow rates greater or lesser than prescribed level.

Nursing-Gen_Chapter 08.indd 144 07/05/18 12:56 PM