Sl.No	NURSING ACTION	RATIONALE
4	Pad the metal frame that supports the canopy.	Protects the child from injury.
5	Flush the tent with oxygen (increase the flow rate) after it has been opened for a period of time to increase the concentration of the gas and reset the flow meter to the original level.	Oxygen is circulated in the tent to adjust the concentration.
6	Analyze and record the tent atmosphere every 1-2 hours. Concentration of 30 to 50% can be achieved in well maintained tents.	Concentration varies with the efficiency of the tent, the rate of flow of oxygen, and the frequency with which tent is opened to the outside environment.
7	Maintain a tight fitting canopy whenever possible, provide nursing care through the sleeves or pockets of the tent.	Prevent oxygen leakage and disruption of the tent atmosphere.
8	Check child's temperature routinely.	Moisture accumulation may result in hypothermia.
9	No smoking' sign should be pasted in the unit.	Oxygen helps in combustion.
10	Record the flow rate of oxygen, alteration in flow rate and the child's reaction.	Serves as a communication between staff members.



Oxyhood

Note

1. Oxygen can be administered to babies using oxygen hood (Oxyhood).
2. Oxygen hood is a plastic device, which is kept over the head of the infant. It permits easy access to the child without loss of oxygen. It helps in efficient delivery of oxygen.
3. While placing hood over the head of the child, the edges of the hood should not rub against the child's chin, neck and shoulders.

Nursing-Gen_Chapter 08.indd 147 07/05/18 12:56 PM