
External hemorrhage	Internal hemorrhage
Clean the area with clean cloth	Do not give anything by mouth.
Place a sterile gauze pad or clean cloth and apply firm pressure at the point to stop bleeding,	If the casuality is unconscious check ABCs
If the bleeding does not stop in 10 minutes elevate the part above heart level to reduce blood flow,	If unconscious put in side lying position to prevent aspiration of secretion.
Reassess circulation every 20-30 minutes. Keep the casuality warm.	Treat the shock by raising the legs 8-12 inches.
Treat the shock by raising the legs 8-12 inches.	Transport immediately for medical care.

Effects of Extreme Heat Stroke

- It occurs when body can no longer controls its temperature anymore and the body's temperature increases to106 degree or higher within 10 to 15 minutes.
- It is caused by very high environment temperature or illness like malaria.
- Exposure to heat and humidity for long time
- Prolonged confinement in hot atmosphere.
- Consumption of alcohol.




Management

- Move the causality to cold place and remove the clothing.
- If the casualty is conscious, then place him in half sitting position with head and shoulders supported.
- If the casualty is unconscious, then place in recovery position.
- Wrap the casualty in a wet sheet and keep it wet. Fan should be on.pour water all over the body. Colds ponging should be started
- Replace the body fluids. Give cold water to drink
- Apply ice cap with ice pieces over the head and neck.
- Cold water enema can be given
- If required, shift him to hospital.



Heat stroke if not treated immediately can cause permanent brain damage.



Have the person lie down

Nursing-Gen_Chapter 09.indd 173 07/05/18 1:01 PM