Severe internal or external bleeding Loss of body fluids, from dehydration, diarrhea, vomiting or burns Severe allergic reactions and severe infection Severe or extensive injuries Electric shock Exposure to extreme heat and cold Bites or stings Gas poisoning Emotional stress, illness.	cold, clammy skin faintness/ dizziness Blood pressure falls Pupils are dilated Lustreless eyes Shaking and trembling of arms and legs Unconsciousness may develop.	Weakness and fainting Giddiness & disorientation Shallow, rapid or gasping breathing Skin become pale, cold and clammy Nausea, vomiting or extreme thirst.

Effects of shock

- Early loss of consciousness
- Failing heart output and insufficient oxygen to cells that are vital for survival.
- Sustained lowered blood pressure

Types of Shock	
Neurogenic Shock	From damage to the nervous system such as a severed spine or a brain injury.
Haemorrhagic Shock	Loss of blood due to wound and internal bleeding.
Respiratory Shock	Insufficient amount of oxygen in the blood due to inadequate breathing.
Cardiogenic Shock	Cardiac muscle not pumping effectively due to injury or previous heart attack.
Circulatory shock,	It is a life-threatening medical condition of low blood perfusion to tissues resulting in cellular injury and inadequate tissue function.
Hypovolemic shock	Related to low blood-volume from hemorrhage, internal bleeding, severe dehydration, vomiting, diarrhea, uncontrolled diabetes, large areas of severely-burned skin, or extreme heat stress.
Metabolic	Loss of body fluids with a change in biochemical equilibrium
Septic Shock	Toxin causes pooling of blood in capillaries not enough blood available for tissues.
Anaphylactic Shock	Severe allergic reaction of the body to sensitization by a foreign protein caused by an allergic reaction to a food, drug, or venom.

Nursing-Gen_Chapter 09.indd 175 07/05/18 1:01 PM