Types of Shock	
Traumatic shock	Brought on by either a traumatic physical injury such as being hit by a car or a mental/emotional blow such as seeing a loved one killed.
Insulin shock	Hypoglycemic (low blood sugar) reaction to an overdose of insulin, a skipped meal, or strenuous exercise.
Electric shock	Injuries caused when electric current passes through body.



Dropping blood pressure to dangerously low levels by widening blood vessels, causing respiratory distress, liver and kidney failure, and coma. Includes **toxic shock syndrome.**

Management

a. Immediately reassure and comfort the casuality.

b. Normally the lower extremities should be elevated. This reduces the blood in the extremities and improves the blood supply to the heart.

c. If there are indications of the head injuries, the head could be raised slightly to reduce pressure on the brain.

d. If there are breathing difficulties, the victim may be more comfortable with head and shoulders raised

e. Loosen the tight clothing to help the circulation and assist breathing.

f. Treat the cause of shock, stop bleeding, immobilize fracture.

g. If breathing and heart beat stop then;-

h. Establish the airway

i. Begin resuscitation immediately.

j. Keep patient in recovery position.

Follow DRSABCD and manage other severe injuries

D - Danger — Ensure the area is safe.

R -Response — Check for response: ask their name, squeeze their shoulders.

S-Send for help

A – Airway — Open patient's mouth.

B-Breathing — Check for breathing: look, listen and feel.

C- CPR — Start CPR: 30 chest compressions followed by 2 breaths.

D- Defibrillation

9.6 Fracture

A break or crack in a bone is called a fracture.

A dislocation is where a bone has been displaced from its normal position at a joint. A fracture is when a bone has been broken.

Nursing-Gen_Chapter 09.indd 176 07/05/18 1:01 PM