Fracture

closed fracture the skin surface around the damaged bone is not broken".

open fracture the skin over the fracture has been damaged or broken.




FLFPDX

Causes;-

Direct force:- A bone may break at the point where a heavy blow is received.

Indirect force:- a bone breaks away from the spot of application of force.

Types of Fracture	
Simple fracture	This is the clean break or crack in the bone
Compound fracture	in this type of fracture the skin is torn by the broken bone which may protrude through the wound
Greenstick fracture	Is a break that happens on oneside of a bone.
Comminuted fracture	This type of fracture produce multiple fragments
Impacted fracture	This type of fracture the ends of the broken bone driven in to each other.



DO YOU KNOW?

Green stick fractures occur in children younger than 10 years because their bones are softer and more flexible than the bones of adults

What to look for – fractures

Swelling.

Pain at or near the site of injury.

Difficulty moving.

Movement in an unnatural direction.

A limb that looks shorter, twisted or bent.

A grating noise or feeling.

Loss of strength .

The aims of first aid for the fractures are:

1. To prevent further damage
2. To reduce pain and shock
3. To make the patient feel comfortable
4. To get medical aid as soon as possible.

If you suspect that someone has fractured a bone

Do's

- Stop any bleeding .
- Immobilize the injured part.
- Apply ice pack wraped in a towel .
- Treat for shock.

Don'ts

- Do not force anyone to use a painful body part.
- Do not straighten a misshapen bone.
- Do not place ice/cold pack directly on skin.
- Do not move victim if neck or spine injury is suspected, unless absolutely necessary.

Nursing-Gen_Chapter 09.indd 177 07/05/18 1:01 PM