- Do not move until injury has been immobilized.
- Do not remove shoes, boots, or clothes around a possible fracture.
- Splinting is unnecessary if victim can give the broken bone sufficient support and immobility.
- Do not splint a possible fractured bone if doing so causes pain.
- Do not Massage the affected area

Points to Remember
DO NOT move the casuality until the injured part is secured and supported unless he/she is in danger
DO NOT let the casuality eat or drink
DO NOT try to replace a disclosed bone into its socket

9.6.1 Dislocation of Joints

A joint is where two bones join or connect. A dislocated joint happens when bones are partly or completely pulled out of their normal position.

The most common joints that dislocate are the shoulder, knee, jaw, or joints in the thumbs or fingers.

Signs of Dislocated joints

The four signs of a dislocated joint are:

1. Strong, sickening pain
2. Not being able to move the joint
3. Swelling and bruising around the joint
4. Shortening, bending or deformity of the joint

First aid for dislocated joints

- Advise them to stay still and help them to support their dislocated joint in the most comfortable position.
- Stop the joint from moving using a bandage. For an arm injury, make a sling to support the arm. For a leg injury, use padding or broad-fold bandages..
- Apply an ice pack. Ice can ease swelling and pain in and around the joint.
- Once you've stopped the joint from moving, take or send the injured person to hospital.
- Keep checking their breathing, pulse and level of response. Check the circulation beyond the bandages every ten minutes and loosen if necessary.

9.7 Insect Bites

Bee, Wasp, Ant stings and other Insect bites.

Remove the sting by scraping, never squeeze the site.

Wash the area and apply antiseptic cream.

Keep the sting site rested, elevated and cool.

To relieve pain and swelling apply cold compress.

Local swelling and irritation may last for several days.

Scorpions Bites

Scorpion stings can be very painful and the pain may persist for several hours. Local redness and numbness often occur.

Nursing-Gen_Chapter 09.indd 178 07/05/18 1:01 PM