




Step 3: Give 30 chest compressions

- Kneel next to the person.
- Place the heel of one hand in the centre of their chest. Place your other hand on top of the first. Interlock your fingers.
- With straight arms, use the heel of your hand to push the breastbone down firmly and smoothly, so that the chest is pressed down between 5–6 cm, and release.
- Do this at a rate of 100 to 120 chest compressions per minute – that's around 2 per second.
- Give **30 chest compressions.**

Step 4: Give two rescue breaths

- Open the airway Place one hand on the person's forehead, gently tilt their head back, then lift their chin using two fingers of your other hand under their chin – when you do this you open their airway.
  - Take a normal breath, make a seal around their mouth and breathe out steadily.
  - The person's chest should rise and fall. Keeping the person's head back and the chin lifted, take your mouth away, take another normal breath, and give a second rescue breath. The two breaths should take no longer than five seconds.



Step 5: Repeat until an ambulance arrives

Repeat 30 compressions and two rescue breaths.

Complications of CPR

- Fractures of ribs or the sternum from chest compression.
- Gastric insufflations from excessive artificial respiration.

9.9 Disaster

A disaster is a sudden calamitous event that seriously disrupts the functioning of a community or society and causes losses that exceeds the community or societys ability to cope using its own resources.

Red Cross (1975) defines Disaster as "An occurrence such as hurricane, tornado, storm, flood, high water, wind-driven water, tidal wave, earthquake, drought, blizzard, pestilence, famine, fire, explosion, building collapse, transportation wreck, or other situation that causes human suffering or creates human that the victims cannot alleviate without assistance."

Nursing-Gen_Chapter 09.indd 181 07/05/18 1:01 PM