- Do not use extra turns in order to use all the bandages.
- When completed, fix the bandage with a circular turn and secure it with a safety pin or other suitable materials such as adhesive strapping.

Preparation of the patient

- Explain the sequence of the procedure to the patient and explain how the patient can assist you.
- Place the articles needed conveniently in the bed side table.
- Bring the patient to the edge of the bed.
- Provide privacy.
- Help the patient to assume comfortable and correct position.
- Perform hand hygiene.

Rules for Application

- Face the patient.
- When bandaging left limb, hold the head of the bandage in the right hand vice versa.
- Apply the outer side of the bandage over the pad and wind it around the injury twice so that it is firm.
- Bandage from below upwards over the limb. Also make it a roll to apply bandage from the inner side to the outer side.
- See that the bandage is neither too loose nor too tight.
- Roll bandage so that each layer covers two-thirds of the earlier layer. Fix the bandage by pinning it up or using adhesive plaster. The usual practice of tearing the final end into two long tails and tying them up is quite satisfactory.

Articals Required

1. Correct width and number of bandages.
2. Disposable gloves (if necessary)
3. Safety pins
4. Scissors
5. Adhesive tapes
6. Rubber Sheet (if necessary)

Procedure

- Apply bandage from distal point toward proximal boundary using variety to turns to cover various shapes of body parts.
- Unroll and very slightly stretch bandage
- Over lap turns by one half to two thirds width of bandage rolls.
- Apply additional rolls without leaving any uncovered skin surface. Secure last bandage applied.
- Remove gloves if worn and perform hand hygiene.
- Assess distal circulation when bandage application is complete and atleast twice during 8 hours period.
- observe the bandage site for 5 – P

It comes in various widths lengths and types of material. For best results, use different widths for different body areas.

For e.g

Fingers	—	1 inches
Hand & arm	—	2 to 2.5 inches
Leg	—	3 to 3.5 inches
Trunk	—	4 to 6 inches

Five 'P'

- Pain
- Pallor
- Pulselessness
- Palpate skin for warmth
- Paralysis

Nursing-Gen_Practical_Unit 10_304-308.indd 307 07/05/18 4:06 PM