wiper from above downwards towards the bulb(to prevent bacteria from spreading all over the thermometer)

7) Collect the dirty soapy water in the kidney tray and place the dirty wiper in the paper bag.
8) Replace thermometer in the test tube of bottle with the lotion
9) Record the temperature in the chart

After care of the equipment

1) Clean all the articles used
2) Wash the thermometer with soap and cold water
3) Keep the thermometer in the antiseptic lotion for 2 to minutes
4) Reset the tray and keep it ready for the next use.

Conversion of Scales

To convert centigrade to Fahrenheit, multiply by 9, divide by 5 and add 32.

To convert Fahrenheit to centigrade, deduct 32, multiply by 5 and divide by 9

$$C = (F-32) \times 5/9$$
$$F = 9/5 \times C+32$$

Abnormal temperature

1) Subnormal - 35°to 36.1°C (95°to 97°F)
2) Collapse – below 35°C (95°F)
3) Fever or pyrexia

Low pyrexia – 37.2° to 38.3°C (99° to 101°F)

Moderate Pyrexia – 38.3° to 39.4°C (101° to 103°F)

High Pyrexia – 39.5°to 40.5°C (103°to 105°F)

Hyper Pyrexia – 40.5°C (105°F) and above



Galileo Galilei was the inventer of Primitive themometer.

5.6 Pulse

Pulse is the expansion felt in an artery where it can be pressed against a bone

Sites

Radial: It is felt with two or three fingers lightly placed on the thumb side of the wrist, anterior surface.

Carotid: This is felt in the neck beside the larynx.

Temporal: It is felt in front of the ear.

Facial: Near the angle of the lower jaw.

Femoral: Felt in the groin

Anterior Fontanel: An infant's pulse may be felt at the 'soft spot' on his head

Observation of the pulse

1. Rate – Number of pulse beat per minute
2. Rhythm or Regularity
3. Strength

Pulse rate

The normal pulse rates for persons at rest are

Adults : 60 to 80 beats/minute

Children : 80 to 100 beats /minute

Infants : 100 to 140 beats /minute

Nursing-Gen_Chapter 05.indd 69 07/05/18 12:48 PM