# NATIONAL IMMUNIZATION SCHEDULE

| National Immunization Schedule for Infants, Children and Pregnant Women | | | | |
|---|---|---|---|---|
| **Vaccine** | **When to give** | **Dose** | **Route** | **Site** |
| **For Pregnant Women** | | | | |
| TT - 1 | Early in pregnancy | 0.5ml | Intra-muscular | Upper Arm |
| TT - 2 | 4 weeks after TT - 1* | 0.5ml | Intra-muscular | Upper Arm |
| TT - Booster | If received 2 TT doses in a pregnancy within last 3 years* | 0.5ml | Intra-muscular | Upper Arm |
| **For Infants** | | | | |
| BCG | At birth or as early as possible till one year of | 0.1ml (0.05ml till 1 month age) | Intra-dermal | Left Upper Arm |
| Hepatitis B | At birth or as early as possible within 24 hours | 0.5ml | Intra-muscular | Antero-lateral side of mid thigh |
| OPV - O | At birth or as early as possible within the first 15 | 2 drops | Oral | Oral |
| OPV 1, 2 & 3 | At 6 weeks, 10 weeks & 14 weeks\ | 2 drops | Oral | Oral |
| DPT 1, 2 & 3 | At 6 weeks, 10 weeks & 14 weeks\ | 0.5ml | Intra-muscular | Antero-lateral side of mid thigh |
| Hep B 1, 2 & 3 | At 6 weeks, 10 weeks & 14 weeks\ | 0.5ml | Intra-muscular | Antero-lateral side of mid thigh |
| Measles | 9 completed months - 12 months | 0.5ml | Sub-cutaneous | Right upper arm |
| Vitamin-A (1st dose) | At 9 months with measles | 1 ml (1 lakh IU) | oral | Oral |
| **For Children** | | | | |
| DPT Booster | 16-24 months | 0.5ml | Intra-muscular | Antero-lateral side of mid thigh |
| Measles 2nd dose | 16-24 months | 0.5ml | Sub-cutaneous | Right upper arm |
| OPV Booster | 16-24 months | 2 drops | Oral | Oral |
| Japanese Encephalitis** | 16-24 months | 0.5ml | Sub-cutaneous | Left Upper Arm |
| Vitamin-A*** | | | | |
| (2nd to 9th dose) | 16 months. Then one dose every 6 months upto the | 2ml (2 lakh IU) | Oral | Oral |
| DPT Booster | 5-6 years | 0.5ml | Intra-muscular | Upper Arm |
| TT | 10 years & 16 years | 0.5ml | Intra-muscular | Upper Arm |

* Give TT-2 or Booster doses before 36 weeks of pregnancy. However, give these even if more than 36 weeks have passed. Give TT to a woman in labour, if she has not previously received TT.

** JE Vaccine, in select endemic districts after the campaign.

*** The 2nd to 9th doses of Vitamin A can be administered to children 1 - 5 years old during biannual rounds, in collaboration with ICDS.



Nursing-Gen_Chapter 06.indd 85 07/05/18 12:51 PM