


	A	B	C	D	E	F	G	H
1	Reg. No	Name	Tam	Eng	CS	Com	Acc	
2	12001	Jayashree	147	136	105	163	162	
3	12002	Kowsalya	156	148	149	147	179	
4	12003	Muskan S	149	165	123	168	179	
5	12004	Ashia Step	168	144	146	192	167	
6	12005	Vennila T I	199	198	150	200	200	
7	12006	Deepika M	187	141	98	130	178	
8	12007	Tharani J	165	102	100	192	192	
9	12008	Thulasi A	143	169	88	176	173	
10	12009	Ayisha B	120	138	109	182	167	
11	12010	Jenifer A	145	135	95	180	185	
12								

Figure 7.22 Worksheet Illustration

7.5.2 Construction of formula

To construct a formula, follow the steps below:

- Cell pointer should be in the cell in which you want to display the result.
- Formula should begin with an = sign.
- In a formula, use only cell reference (cell addresses) instead of the actual values within the cells.
- While constructing a formula, ***BODMAS rule*** should be kept in mind.
- General Syntax of constructing a formula is: = ***cell reference1 <operator> cell reference2 <operator> .....***
- Cell references are of two types (i) Relative cell reference (ii) Absolute Cell reference.
- If you refer cell addresses directly while constructing formulae, it is called as "Relative Cell reference".
- Examples of Relative Cell references:

Adding values of A1, B1, C1, D1	=A1+B1+C1+D1
Subtract E4 from H3	= H3 – E4
Multiply A5 and B5	= A5 * B5
Average of G1, G2, G3, G4	=(G1+G2+G3+G4)/4

Chapter 7.indd 117 3/20/2020 11:37:10 AM