- In the above table, all cell references are “Relative cell references”.
- While writing a formula, if you use the $ symbol in front of a column name and row number, it will become an “Absolute Cell reference”.
- Examples of Absolute cell references:

Adding values of A1, B1, C1, D1	=$A$1+$B$1+$C$1+$D$1
Subtract E4 from H3	= $H$3 – $E$4
Multiply A5 and B5	= $A$5 * B5
Average of G1, G2, G3, G4	=($G$1+G2+$G$3+G4)/4

- In an expression, all cells need not necessarily be relative or absolute. You can mix both type of references.
- The following section explains the use of relative cell reference. About “Absolute cell reference”, you will be learn later in this chapter.

Finding Total to the above Illustration:

- Move the cell pointer to H2 (Total column)
- Type the following formula; after entering the formula, press “Enter” key

  = C2+D2+E2+F2+G2 (Refere Figure 7.25)

- Now, you will get the sum of all the values of C2, D2, E2, F2 and G2
- The above-mentioned formula clearly stated that, how worksheets are working with cells.
- While referring to the cell addresses in a formula, the spreadsheet reads the value inside the cell that you refer. This is a good practice of constructing a formula. Because, if you change any value, the spreadsheet recalculates with that new value.

  After entering a formula the result is display as in Figure 7.23

Chapter 7.indd 118 3/20/2020 11:37:10 AM