var m = n = z = 25; // sets all three variables to a value of 25//

The assignment operator can also be used to set a variable to hold the value of an expression. For example,

var x = 102 + 5 - 50; // x set to 57 //

JavaScript supports some shorthand arithmetic operators like +=, -=, *=, /= and %= to evaluate arithmetic calculations.

Table: 14.2 Shorthand Arithmetic operators

Shorthand Arithmetic Operator	Meaning	Example	Result
+=	Add and assign	var sum = 120; sum += 20;	Variable sum = 140
-=	Subtract and assign	var diff = 120; diff –= 20;	Variable diff = 100
*=	Multiply and assign	var prod = 100; prod *=10;	Variable prod = 1000
/=	Division	Var res = 522; Res/=100	Variable res = 5.22
%=	Modulus operator	Var rem = 522; rem %= 100	Variable rem = 22 (remainder)

Illustration 14.4 Using Arithmetic Shorthand Operators

```
<Html>
        <Head>
                <Title>Demo Program - To test Arithmetic Shorthand Operators in JavaScript
</Title>
        </Head>
<Html>
        <Head>
                <Title>Demo Program - To test Arithmetic Shorthand Operators in JavaScript
</Title>
        </Head>
<Body>
        <script language="javascript" type="text/javascript">
        var value1 = 522, value2=10;
        document.write("<br>Data1 : "+value1);
        document.write("<br>Data2 : "+value2);
        var sum = value1; sum+=value2;
        var diff = value1; diff-=value2;
        var prod = value1; prod*=value2;
        var res = value1; res/=value2;
```


Chapter 14 Pages 127-153.indd 301 3/20/2020 11:44:36 AM