33. Following data represents the life of two models of refrigerators A and B.

Life (No. of years)		0-2	2-4	4-6	6-8	8-10	10-12
Refrigerator	Model A	5	16	13	7	5	4
	Model B	2	7	12	19	9	1

Find the mean life of each model. Which model has greater uniformity? Also obtain mode for both models.

34. Calculate the quartile deviation and coefficient of quartile deviation from the following data:

Size	06	09	12	15	18
Frequency	7	12	19	10	2

35. Calculate the appropriate measure of dispersion from the following data:

Wages in ₹	Below 35	35-37	38-40	41-43	Above 43
Earners	14	60	95	24	7

36. Two brands of tyres are tested with the following results:

Life (in 1000 miles)	20-25	25-30	30-35	35-40	40-45	45-50
Brand A	8	15	12	18	13	9
Brand B	6	20	32	30	12	0

Which is more consistent ?

37. Find the S.D. for the number of days patients admitted in a hospital.

Days of confinement	5	6	7	8	9
No. of patients	18	14	9	3	1

38. Calculate the quartile deviation and its coefficient from the following data:

Class Interval	10 -15	15 - 20	20 - 25	25 - 30	30 - 35
Frequency	8	12	14	10	6

Ch6_11th Stats.indd 187 2/3/2019 11:02:22 AM