39. Calculate the Mean Deviation from mean and its Coefficient from the following data, relating to Height (to the nearest cm) of 100 children:

| Height(cms) | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 |
|---|---|---|---|---|---|---|---|---|---|
| No. of Children | 2 | 0 | 15 | 29 | 25 | 12 | 10 | 4 | 3 |

40. Find the standard deviation for the distribution given below:

| x | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
|---|---|---|---|---|---|---|---|
| Frequency | 10 | 20 | 30 | 35 | 14 | 10 | 2 |

41. Calculate the coefficient of range separately for the two sets of data:

| Set 1 | 10 | 20 | 9 | 15 | 10 | 13 | 28 |
|---|---|---|---|---|---|---|---|
| Set 2 | 35 | 42 | 50 | 32 | 49 | 39 | 33 |

42. Blood serum cholesterol levels of 10 persons are 240, 260, 290, 245, 255, 288, 272, 263, 277 and 250. Calculate the standard deviation with the help of assumed mean.

43. Two groups of people played a game which reveals the quick operation of a particular key in a computer and the fraction of reaction times nearest to the tenth of a second is given below .

| | Minimum | I Quartile | Median | III Quartile | Maximum |
|---|---|---|---|---|---|
| Group I | 0.6 | 0.8 | 1.0 | 1.5 | 1.9 |
| Group II | 0.4 | 0.7 | 1.0 | 1.3 | 1.6 |

Draw two Box-Whisker plots and compare reaction times of the two groups.

44. Draw Box – Whisker plot for the following

(i) 3, 5, 10, 11, 12, 16, 17, 17, 19, 20, 22

(ii) –7, –5, –4, –4, –3, –3, –2, –1, 0, 1, 4, 6, 8, 9

**Answers**

I.1. d 2. b 3. c 4. d 5. a 6. c 7. a 8. c 9. d 10. c

II. 11. IQR 12. variance, standard deviation 13.coefficient of variation

14. 50% 15. normal (or) symmetric V.29. 2nd Formula 30. typist B

31. 7.13 32. 51.57, 7.67 33. (i) 5.12, 6.16 (ii) model B

34. 3, 0.25 35. $QD$=1.8, $CQD$=0.047 36. brand B 37. 1.03 38. 4.94,0.2265

39. 1.24, 0.0194 40. 1.4 41. 0.51,0.22 42. 16.5



Ch6_11th Stats.indd 188 2/3/2019 11:02:22 AM