- The column or row heads should indicate the units of measurements such as monetary units like Rupees, and other units such as meters, etc. wherever necessary.
- Column heading may be numbered for comparison purposes. Items may be arranged either in the order of their magnitude or in alphabetical, geographical, and chronological or in any other suitable arrangement for meaningful presentation.
- Figures as accurate as possible are to be entered in a table. If the figures are approximate, the same may be properly indicated.

3.6 Frequency Distribution

A tabular arrangement of raw data by a certain number of classes and the number of items (called frequency) belonging to each class is termed as a frequency distribution. The frequency distributions are of two types, namely, discrete frequency distribution and continuous frequency distribution.

3.6.1 Discrete Frequency Distribution

Raw data sometimes may contain a limited number of values and each of them appeared many numbers of times. Such data may be organized in a tabular form termed as a simple frequency distribution. Thus the tabular arrangement of the data values along with the frequencies is a simple frequency distribution. A simple frequency distribution is formed using a tool called 'tally chart'. A tally chart is constructed using the following method:

- Examine each data value.
- Record the occurrence of the value with the slash symbol (/), called tally bar or tally mark.
- If the tally marks are more than four, put a crossbar on the four tally bar and make this as block of 5 tally bars (𝍸)
- Find the frequency of the data value as the total number of tally bars i.e., tally marks corresponding to that value.

Example 3.11

The marks obtained by 25 students in a test are given as follows: 10, 20, 20, 30, 40, 25, 25, 30, 40, 20, 25, 25, 50, 15, 25, 30, 40, 50, 40, 50, 30, 25, 25, 15 and 40. The following discrete frequency distribution represents the given data:

Ch3_11th Stats.indd 54 2/3/2019 10:58:16 AM