Table 3.13(ii)

Marks secured by students (Exclusive Method)

Marks	x, Integer value	Tally Marks	No. of Students
20-25	$20 \leq x < 25$	𝍸	5
25-30	$25 \leq x < 30$	𝍸 //	7
30-35	$30 \leq x < 35$	𝍸 ////	9
35-40	$35 \leq x < 40$	𝍸 𝍸 /	11
40-45	$40 \leq x < 45$	𝍸 𝍸 //	12
45-50	$45 \leq x < 50$	𝍸 /	6
		Total	**50**

True class intervals

In the case of continuous variables, we take the classes in such a way that there is no gap between successive classes. The classes are defined in such a way that the upper limit of each class is equal to lower limit of the succeeding class. Such classes are known as true classes. The inclusive method of forming class intervals are also known as not-true classes. We can convert the not-true classes into true-classes by subtracting 0.5 from the lower limit of the class and adding 0.5 to the upper limit of each class like 19.5 - 25.5, 25.5 - 30.5, 30.5 – 35.5, 35.5 – 40.5, 40.5 - 45.5, 45.5 – 50.5.

Open End Classes

When a class limit is missing either at the lower end of the first class interval or at the upper end of the last classes or when the limits are not specified at both the ends, the frequency distribution is said to be the frequency distribution with open end classes.

Example 3.14

Salary received by 113 workers in a factory are classified into 6 classes. The classes and their frequencies are displayed in Table 3.14 Since the lower limit of the first class and the upper limit of the last class are not specified, they are open end classes.

Table 3.14

Open-Ended Frequency Table

Salary Range in Rs.	Number of workers
Below 10000	18
10000 - 20000	23
20000 - 30000	30

Ch3_11th Stats.indd 58 2/3/2019 10:58:16 AM