



Lysome

Centrioles


They are generally found close to the nucleus and are made up of tube-like structures. Centrioles or centrosomes are present only in animal cells and absent in plant cells. It helps in the separation of chromosomes during cell division.



Structure of a Centriole

Endoplasmic reticulum - You guys, be quiet, I have so much work to do

It is an inter membranous network made up of flat or tubular sacs within the cytoplasm. Endoplasmic reticulum is of two types. They are rough endoplasmic reticulum and smooth endoplasmic reticulum.



Endoplasmic reticulum

Rough endoplasmic reticulum are rough due to the ribosomes attached to the membrane. which helps in the synthesis of protein.


Smooth endoplasmic reticulum. It is a network of tubular sacs without ribosomes on the membrane. They play a role in the synthesis of lipids, steroids and also transport them within the cell.

Nucleus - Everyone do what I say. Acting like the "Brain" of the cell

- Plant and animal cells have a nucleus inside the cytoplasm. It is surrounded by a nuclear envelope. One or two nucleolus and the chromatin body are present inside the nucleus. During cell division, the chromatin body is organised into a chromosome. Storage of genetic material and transfers heredity characters from generation to generation are the functions of chromosome.

Functions of Nucleus

- In controls all the processes and chemical reactions that take place inside the cell.
- Inheritance of character from one generation to another.



Nucleus

VII Std Science Term-2 EM Unit 4 Folder.indd 66 20-07-2019 1.48.46 PM