with inferior or sometimes with half inferior ovary. Calyciflorae includes 5 orders and 27 families.

**Sub-class 2. Gamopetalae:** Plants with united petals, which are either partially or completely fused to one another and dichlamydeous are placed under Gamopetalae. It is further divided into three series – Inferae, Heteromerae and Bicarpellatae.

**Series (i) Inferae:** The flowers are epigynous and with inferior ovary. Inferae includes 3 orders and 9 families.

**Series (ii) Heteromerae:** The flowers are hypogynous, superior ovary and with more than two carpels. Heteromerae includes 3 orders and 12 families.

**Series (iii) Bicarpellatae:** The flowers are hypogynous, superior ovary and with two carpels. Bicarpellatae includes 4 orders and 24 families.

**Sub-class 3. Monochlamydeae:** Plants with incomplete flowers either apetalous or with undifferentiated calyx and corolla are placed under Monochlamydeae. The sepals and petals are not distinguished and they are called **perianth.** Sometimes both the whorls are absent. Monochlamydeae includes 8 series and 36 families.

**Class II Gymnospermae:** Plants that contain naked seeds come under this class. Gymnospermae includes three families – Gnetaceae, Coniferae and Cycadaceae.

**Class III Monocotyledonae:** Plants contain only one cotyledon in their seed, leaves with parallel venation, fibrous root system and trimerous flowers come under this class. The Monocotyledonae has 7 series and 34 families.

The Bentham and Hooker system of classification is still supposed to be the best system of classification. It has been widely practiced in colonial countries and herbaria of those countries were organised based on this system and is still used as a key for the





**Figure 5.6:** Bentham and Hooker system of classification



XIth_Botany_Chapter_5_EM.indd 111 5/17/2020 9:06:00 PM