Amplified Fragment Length Polymorphism (AFLP)

This method is similar to that of identifying RFLPs in that a restriction enzyme is used to cut DNA into numerous smaller pieces, each of which terminates in a characteristic nucleotide sequence due to the action of restriction enzymes.

AFLP is largely used for population genetics studies, but has been used in studies of closely related species and even in some cases, for higher level cladistic analysis.

Random Amplified Polymorphic DNA (RAPD)

It is a method to identify genetic markers using a randomly synthesized primer that will anneal (recombine (DNA) in the double stranded form) to complementary regions located in various locations of isolated DNA. If another complementary site is present on the opposing DNA strand at a distance that is not too great (within the limits of PCR) then the reaction will amplify this region of DNA.

RAPDs like microsatellites may often be used for genetic studies within species but may also be successfully employed in phylogenetic studies to address relationships within a species or between closely related species. However RAPD analysis has the major disadvantage that results are difficult to replicate and in that the homology of similar bands in different taxa may be nuclear.

Significance of Molecular Taxonomy

1. It helps to identify a very large number of species of plants and animals by the use of conserved molecular sequences.
2. Using DNA data evolutionary patterns of biodiversity are now investigated.
3. DNA taxonomy plays a vital role in phytogeography, which ultimately helps in genome mapping and biodiversity conservation.
4. DNA- based molecular markers used for designing DNA based molecular probes, have also been developed under the branch of molecular systematics.

5.11.6 DNA Barcoding

Have you seen how scanners are used in supermarkets to distinguish the **Universal Product Code** (UPC)? In the same way we can also distinguish one species from another. DNA barcoding is a taxonomic method that uses a very short genetic sequence from a standard part of a genome. The genetic sequence used to identify a plant is known as "**DNA tags**" or "**DNA barcodes**". **Paul Hebert** in 2003 proposed '**DNA barcoding**' and he is considered as 'Father of barcoding'.

The gene region that is being used as an effective barcode in plants is present in two genes of the chloroplast, **matK** and **rbcL**, and have been approved as the barcode regions for plants.

Sequence of unknown species can be matched from submitted sequence in GenBank using Blast (web-programme for searching the closely related sequence).

Significance of DNA barcoding

1. DNA barcoding greatly helps in identification and classification of organism.
2. It aids in mapping the extent of biodiversity.

DNA barcoding techniques require a large database of sequences for comparison and prior knowledge of the barcoding region.

However, DNA barcoding is a helpful tool to determine the authenticity of botanical material in whole, cut or powdered form.

XIth_Botany_Chapter_5_EM.indd 117 5/17/2020 9:06:02 PM