Microscopic measurements:

The microscope also has facility to measure microscopic objects through a technique called '**micrometry**'. There are two scales involved for measuring.

1. Ocular Micrometre
2. Stage Micrometre

Ocular Micrometre: It is fixed inside the eye piece lens. It is a thin transparent glass disc where there are lines divided into 100 equal units. The scale has no value.

Stage Micrometre: This is a slide with a line divided into 100 units. The line is about 10mm. The distance between two adjacent lines is 10 µm. The known value of the stage micrometre is transferred to the ocular micrometre, thereby the measurement can be made using ocular micrometre.




Figure 6.5: a. Ocular micrometre; b. Stage micrometre

The distance between two adjacent line of ocular meter= $\frac{\text{Number of stage divisions}}{\text{Number of ocular divisions}} \times 10$

These observations led to the formulation of modern cell theory.

- All organisms are made up of cells.
- New cells are formed by the division of pre-existing cells.
- Cells contains genetic material, which is passed on from parents to daughter cells.
- All metabolic reactions take place inside the cells.

6.3.1 Exception to Cell Theory

Viruses are puzzle in biology. Viruses, viroids and prions are the exception to cell theory. They lack protoplasm, the essential part of the cell and exists as obligate parasites which are sub-cellular in nature.

6.3.2 Protoplasm Theory

Corti first observed protoplasm. **Felix Dujardin** (1835) observed a living juice in animal cell and called it "**Sarcode**". **Purkinje** (1839) coined the term protoplasm for sap inside a plant cell. **Hugo Van Mohl** (1846) indicated importance of protoplasm.

Max Schultze (1861) established similarity between Protoplasm and Sarcode and proposed a theory which later on called "**Protoplasm Theory**" by **O. Hertwig** (1892). **Huxley** (1868) proposed Protoplasm as a "**physical basis of life**".

Protoplasm as a Colloidal System

Protoplasm is a complex colloidal system which was suggested by **Fisher** in 1894 and **Hardy** in 1899. It is primarily made of water and various other solutes of biological importance such as glucose, fatty acids, amino acids, minerals, vitamins, hormones and enzymes.

These solutes may be homogeneous (soluble in water) or heterogeneous mass (insoluble in water) which forms the basis for its colloidal nature.

Physical Properties of Protoplasm

The protoplasm exists either in semisolid (jelly-like) state called '**gel**' due to suspended particles and various chemical bonds or may be liquid state called '**sol**'.The colloidal

XIth_Botany_Chapter_6_EM.indd 147 5/17/2020 9:09:43 PM