Fluid Mosaic Model

Jonathan Singer and **Garth Nicolson** (1972) proposed fluid mosaic model.

It is made up of lipids and proteins together with a little amount of carbohydrate. The lipid membrane is made up of phospholipid. The phospholipid molecule has a hydrophobic tail and hydrophilic head. The hydrophobic tail repels water and hydrophilic head attracts water. The proteins of the membrane are globular proteins which are found intermingled between the lipid bilayer most of which are projecting beyond the lipid bilayer. These proteins are called as **integral proteins**. Few are superficially attached on either surface of the lipid bilayer which are called as **peripheral proteins**. The proteins are involved in transport of molecules across the membranes and also act as enzymes, receptors (or) antigens.



Water-loving polar molecule are called hydrophilic molecule. They have polar phosphate group responsible for attracting water.

Water hating non-polar molecule are called as hydrophobic molecule. They have fatty acid which is non-polar which cannot attract water

Carbohydrate molecules of cell membrane are short chain polysaccharides. These are either bound with '**glycoproteins**' or '**glycolipids**' and form a '**glyocalyx**' (Figure 6.11).

The movement of membrane lipids from one side of the membrane to the other side by vertical movement is called **flip flopping** or **flip flop movement**. This movement takes place more slowly than lateral diffusion of lipid molecule. The Phospholipids can have flip flop movement because they have smaller polar regions, whereas the proteins cannot flip flop because the polar region is extensive.

Function of Cell Membrane

The functions of the cell membrane is enormous which includes cell signalling, transporting nutrients and water, preventing unwanted substances entering into the cell, and so on.

Cytoplasm

Cytoplasm is the main arena of various activities of a cell. It is the semifluid gelatinous substance that fills the cell. It is made up of eighty percent water and is usually clear and colourless. The cytoplasm is sometimes described as non nuclear content of protoplasm. The cytoplasm serves as a molecular soup where all the cellular organelles are suspended and bound together by a lipid bilayer plasma membrane. It constitutes dissolved nutrients, numerous salts and acids to dissolve waste products. It is a very good conductor of electricity. It gives support and protection to the cell organelles. It helps movement of the cellular materials around the cell through a process called **cytoplasmic streaming**. Further, most cellular activities such as many metabolic pathways including glycolysis and cell division occur in cytoplasm.

6.6 Cell Organelles

6.6.1 Endomembrane System

System of membranes in a eukaryotic cell, comprises the plasma membrane, nuclear membrane, endoplasmic reticulum, golgi apparatus, lysosomes and vacuolar membranes (tonoplast). Endomembranes are made up of phospholipids with embedded proteins that are similar to cell membrane which occur within the cytoplasm. The endomembrane system is evolved from the inward growth of cell membrane in the ancestors of the first eukaryotes (Figure 6.12).

6.6.2 Endoplasmic Reticulum

The largest of the internal membranes is called the **endoplasmic reticulum** (ER). The name endoplasmic reticulum was given by **K.R. Porter** (1948). It consists of double membrane. Morphologically the structure of endoplasmic reticulum consists of the following:

XIth_Botany_Chapter_6_EM.indd 153 5/17/2020 9:09:51 PM