- Secretion of carbohydrates for the formation of plant cell walls and insect cuticles.
- **Zymogen granules** (proenzyme/precursor of all enzyme) are synthesised.




Figure 6.14: Exocytosis

6.6.4 Mitochondria

It was first observed by **A. Kolliker** (1880). **Altmann** (1894) named it as Bioplasts. Later **Benda** (1897, 1898), named as mitochondria. They are ovoid, rounded, rod shape and pleomorphic structures. Mitochondrion consists of double membrane, the outer and inner membrane. The outer membrane is smooth, highly permeable to small molecules and it contains proteins called **Porins**, which form channels that allows free diffusion of molecules smaller than about 1000 daltons and the inner membrane divides mitochondrion into two compartments, outer chamber between two membranes and the inner chamber is filled with matrix.

The inner membrane is convoluted (infoldings), called **crista** (plural: cristae). Cristae contain most of the enzymes for electron transport system. Inner chamber of the mitochondrion is filled with proteinaceous material called **mitochondrial matrix**. The Inner membrane consists of stalked particles called **elementary particles** or **Fernandez Moran particles**, **F1 particles** or **Oxysomes.** Each particle consists of a base, stem and a round head. In the head, ATP synthase is present for oxidative phosphorylation. Inner membrane is impermeable to most ions, small molecules and maintains the proton gradient that drives oxidative phosphorylation (Figure 6.15).




Figure 6.15: Structure of Mitochondria

Mitochondria contain 73% of proteins, 25-30% of lipids, 5-7 % of RNA, DNA (in traces) and enzymes (about 60 types). Mitochondria are called **Power house of a cell**, as they produce energy rich ATP.

All the enzymes of Kreb's cycle are found in the matrix except succinate dehydrogenase. Mitochondria consist of circular DNA and 70S ribosome. They multiply by fission and replicates by strand displacement model. Because of the presence of DNAs it is semi-autonomous organelle. Unique characteristic of mitochondria is that they are inherited from female parent only. Mitochondrial DNA comparisons are used to trace human origins. It is also used to track and date recent evolutionary time because it mutates 5 to 10 time faster than DNA in the nucleus.

6.6.5 Plastids

The term plastid is derived from the Greek word ***Platikas*** (formed/moulded) and used by **A.F.U. Schimper** in 1885. He classified plastids into following types according to their structure, pigments and function. Plastids multiply by fission.

XIth_Botany_Chapter_6_EM.indd 155 5/17/2020 9:09:52 PM