They contain a variety of hydrolytic enzymes, that can digest material within the cell. The membrane around lysosome prevent these enzymes from digesting the cell itself (Figure 6.18).




Figure 6.18: Structure of Lysosome

Functions:

- **Intracellular digestion:** They digest carbohydrates, proteins and lipids present in cytoplasm.
- **Autophagy:** During adverse condition they digest their own cell organelles like mitochondria and endoplasmic reticulum
- **Autolysis:** Lysosome causes self destruction of cell.
- **Ageing:** Lysosomes have autolytic enzymes that disrupts intracellular molecules.




Figure 6.19: Enzymes of Lysosome

- **Phagocytosis:** Large cells or contents are engulfed and digested by macrophages, thus forming a phagosome in cytoplasm. These phagosome fuse with lysosome for further digestion.
- **Exocytosis:** Lysosomes release their enzymes outside the cell to digest other cells (Figure 6.19).

6.6.9 Microbodies

Eukaryotic cells contain many enzyme bearing membrane enclosed vesicles called **microbodies**. They are single unit membrane bound cell organelles. Example: Peroxisomes and glyoxysomes.

6.6.10 Peroxisomes




Figure 6.20: Structure of Peroxisome

Peroxisomes were identified as organelles by **Christian de Duve** (1967). Peroxisomes are small spherical bodies and single membrane bound organelle. It takes part in photorespiration and associated with glycolate metabolism. In plants, leaf cells have many peroxisomes. It is also commonly found in liver and kidney of mammals. These are also found in cells of protozoa and yeast (Figure 6.20).

6.6.11 Glyoxysomes

Glyoxysome was discovered by **Harry Beevers** (1961). It is a single membrane bound organelle. It is a sub cellular organelle and contains enzymes of glyoxylate pathway. β-oxidation of fatty acid occurs in glyoxysomes of germinating seeds Example: Castor seeds.

6.6.12 Sphaerosomes

It is spherical in shape and enclosed by single unit membrane. Example: Storage of fat in the endosperm cells of oil seeds.

6.6.13 Centrioles

Centrioles consists of nine triplet peripheral fibrils made up of tubulin. The central part of the centriole is called **hub**, is connected to the tubules of the peripheral triplets by radial spokes (9+0 pattern). The centriole form the basal body of cilia or flagella and spindle fibers which forms the spindle apparatus in animal

XIth_Botany_Chapter_6_EM.indd 158 5/17/2020 9:09:54 PM