lipids, carbohydrates and cell organelles including mitochondria and endoplasmic reticulum. Many checkpoints control the cell cycle. The check point are also called as the **restriction point.** First check point at the end of G_1, determines a cells fate whether it will continue in the cell cycle and divide or enter a stage called **G_0** a quiescent stage, probably as specified cell or die. Cells are arrested in G_1 due to:

- Nutrient deprivation
- Lack of growth factors or density dependant inhibition
- Undergo metabolic changes and enter into G_0 state.

Biochemicals inside cell activates the cell division. The proteins called **kinases** and cyclins activate genes and their proteins to perform cell division. Cyclins act as major checkpoint which operates in G_1 to determine whether or not a cell divides.



Dolly

Since the DNA of cells in G_0, do not replicate. Researchers are able to fuse the donor cells from a sheep's mammary glands into G_0 state by culturing in the nutrient free state. The G_0 donor nucleus synchronised with cytoplasm of the recipient egg, which developed into the clone Dolly.

7.2.4 G_0 Phase

Some cells exit G_1 and enters a quiescent stage called **G_0**, where the cell remains metabolically active without proliferation. Cells can exist for long periods in G_0 phase. In G_0, cells cease growth with reduced rate of RNA and protein synthesis. The G_0 phase is not permanent. Mature neuron and skeletal muscle cell remain permanently in G_0. Many cells in animals remains in G_0 unless called on to proliferate by appropriate growth factors or other extracellular signals. G_0 cells are not dormant.

7.2.5 S phase – Synthesis phase – cells with intermediate amounts of DNA.

Growth of the cell continues as replication of DNA occur, protein molecules called **histones** are synthesised and attach to the DNA. The centrioles duplicate in the cytoplasm. DNA content increases from 2C to 4C.

7.2.6 G_2 – The second Gap phase – 4C amount of DNA in cells of G_2 and mitosis

Cell growth continues by protein and cell organelle synthesis, mitochondria and chloroplasts divide. DNA content remains as 4C. Tubulin is synthesised and microtubules are formed. Microtubles organise to form spindle fibre. The spindle begins to form and nuclear division follows.

One of the proteins synthesized only in the G_2 period is known as **Maturation Promoting Factor (MPF)**. It brings about condensation of interphase chromosomes into the mitotic form.

DNA damage checkpoints operates in G_1 S and G_2 phases of the cell cycle.

7.3 Cell Division

7.3.1 Amitosis (Direct Cell Division)

Amitosis is also called **direct** or **incipient cell division**. Here there is no spindle formation and chromatin material does not condense. It consist of two steps: (Figure 7.2).




Figure 7.2: Amitosis

XIth_Botany_Chapter_7_EM.indd 168 5/17/2020 9:12:09 PM